

# Algebraic Number Theory

(PARI-GP version 2.15.3)

## Binary Quadratic Forms

create  $ax^2 + bxy + cy^2$  **Qfb**( $a, b, c$ ) or **Qfb**( $[a, b, c]$ )  
reduce  $x$  ( $s = \sqrt{D}$ ,  $l = \lfloor s \rfloor$ ) **qfbred**( $x, \{flag\}, \{D\}, \{l\}, \{s\}$ )  
return  $[y, g]$ ,  $g \in \text{SL}_2(\mathbf{Z})$ ,  $y = g \cdot x$  reduced **qfbreds12**( $x$ )  
composition of forms  $x*y$  or **qfbnucomp**( $x, y, l$ )  
 $n$ -th power of form  $x^n$  or **qfbnupow**( $x, n$ )  
composition **qfbcomp**( $x, y$ )  
... without reduction **qfbcomppraw**( $x, y$ )  
 $n$ -th power **qfbpow**( $x, n$ )  
... without reduction **qfbpowraw**( $x, n$ )  
prime form of disc.  $x$  above prime  $p$  **qfbprimeform**( $x, p$ )  
class number of disc.  $x$  **qfbclassno**( $x$ )  
Hurwitz class number of disc.  $x$  **qfbhclassno**( $x$ )  
solve  $Q(x, y) = n$  in integers **qfbsolve**( $Q, n$ )  
solve  $x^2 + Dy^2 = p$ ,  $p$  prime **qfbcornacchia**( $D, p$ )  
...  $x^2 + Dy^2 = 4p$ ,  $p$  prime **qfbcornacchia**( $D, 4 * p$ )

## Quadratic Fields

quadratic number  $\omega = \sqrt{x}$  or  $(1 + \sqrt{x})/2$  **quadgen**( $x$ )  
minimal polynomial of  $\omega$  **quadpoly**( $x$ )  
discriminant of **Q**( $\sqrt{x}$ ) **quaddisc**( $x$ )  
regulator of real quadratic field **quadregulator**( $x$ )  
fundamental unit in  $O_D$ ,  $D > 0$  **quadunit**( $D, \{ 'w \}$ )  
norm of fundamental unit in  $O_D$  **quadunitnorm**( $D$ )  
index of  $O_{Df_2}^\times$  in  $O_D^\times$  **quadunitindex**( $D, f$ )  
class group of **Q**( $\sqrt{D}$ ) **quadclassunit**( $D, \{flag\}, \{t\}$ )  
Hilbert class field of **Q**( $\sqrt{D}$ ) **quadhilbert**( $D, \{flag\}$ )  
... using specific class invariant ( $D < 0$ ) **polclass**( $D, \{inv\}$ )  
ray class field modulo  $f$  of **Q**( $\sqrt{D}$ ) **quadrays**( $D, f, \{flag\}$ )

## General Number Fields: Initializations

The number field  $K = \mathbf{Q}[X]/(f)$  is given by irreducible  $f \in \mathbf{Q}[X]$ .  
We denote  $\theta = \bar{X}$  the canonical root of  $f$  in  $K$ . A  $nf$  structure contains a maximal order and allows operations on elements and ideals. A  $bnf$  adds class group and units. A  $bnr$  is attached to ray class groups and class field theory. A  $rnf$  is attached to relative extensions  $L/K$ .

init number field structure  $nf$  **nfinit**( $f, \{flag\}$ )  
  known integer basis  $B$  **nfinit**( $[f, B]$ )  
  order maximal at  $vp = [p_1, \dots, p_k]$  **nfinit**( $[f, vp]$ )  
  order maximal at all  $p \leq P$  **nfinit**( $[f, P]$ )  
  certify maximal order **nfcertify**( $nf$ )

### nf members:

a monic  $F \in \mathbf{Z}[X]$  defining  $K$  **nf.pol**  
number of real/complex places **nf.r1/r2/sign**  
discriminant of  $nf$  **nf.disc**  
primes ramified in  $nf$  **nf.p**  
 $T_2$  matrix **nf.t2**  
complex roots of  $F$  **nf.roots**  
integral basis of  $\mathbf{Z}_K$  as powers of  $\theta$  **nf.zk**  
different/codifferent **nf.diff**, **nf.codiff**  
index  $[\mathbf{Z}_K : \mathbf{Z}[X]/(F)]$  **nf.index**  
recompute  $nf$  using current precision **nfnewprec**( $nf$ )  
init relative  $rnf$   $L = K[Y]/(g)$  **rnfinit**( $nf, g$ )  
init  $bnf$  structure **bnfinit**( $f, 1$ )

**bnf members:** same as  $nf$ , plus  
  underlying  $nf$  **bnf.nf**  
  class group, regulator **bnf.clgp**, **bnf.reg**  
  fundamental/torsion units **bnf.fu**, **bnf.tu**  
  add  $S$ -class group and units, yield  $bnfS$  **bnfsunit**( $bnf, S$ )  
  init class field structure  $bnr$  **bnrinit**( $bnf, m, \{flag\}$ )  
**bnr members:** same as  $bnf$ , plus  
  underlying  $bnf$  **bnr.bnf**  
  big ideal structure **bnr.bid**  
  modulus  $m$  **bnr.mod**  
  structure of  $(\mathbf{Z}_K/m)^*$  **bnr.zkst**

## Fields, subfields, embeddings

**Defining polynomials, embeddings**  
(some) number fields with Galois group  $G$  **nflist**( $G$ )  
... and  $|\text{disc}(K)| = N$  and  $s$  complex places **nflist**( $G, N, \{s\}$ )  
... and  $a \leq |\text{disc}(K)| \leq b$  **nflist**( $G, [a, b], \{s\}$ )  
smallest poly defining  $f = 0$  (slow) **polredabs**( $f, \{flag\}$ )  
small poly defining  $f = 0$  (fast) **polredbest**( $f, \{flag\}$ )  
monic integral  $g = Cf(x/L)$  **poltomonic**( $f, \{\&L\}$ )  
random Tschirnhausen transform of  $f$  **poltschirnhaus**( $f$ )  
 $\mathbf{Q}[t]/(f) \subset \mathbf{Q}[t]/(g)$  ? Isomorphic? **nfisincl**( $f, g$ ), **nfisisom**  
reverse polmod  $a = A(t) \bmod T(t)$  **modreverse**( $a$ )  
compositum of  $\mathbf{Q}[t]/(f)$ ,  $\mathbf{Q}[t]/(g)$  **polcompositum**( $f, g, \{flag\}$ )  
compositum of  $K[t]/(f)$ ,  $K[t]/(g)$  **nfcompositum**( $nf, f, g, \{flag\}$ )  
splitting field of  $K$  (degree divides  $d$ ) **nfsplitting**( $nf, \{d\}$ )  
signs of real embeddings of  $x$  **nfeltsign**( $nf, x, \{pl\}$ )  
complex embeddings of  $x$  **nfeltembed**( $nf, x, \{pl\}$ )  
 $T \in K[t]$ , # of real roots of  $\sigma(T) \in R[t]$  **nfpolsturm**( $nf, T, \{pl\}$ )

### Subfields, polynomial factorization

subfields (of degree  $d$ ) of  $nf$  **nfsubfields**( $nf, \{d\}$ )  
maximal subfields of  $nf$  **nfsubfieldsmax**( $nf$ )  
maximal CM subfield of  $nf$  **nfsubfieldscm**( $nf$ )  
 $K_d \subset \mathbf{Q}(\zeta_n)$ , using Gaussian periods **polsubcyclo**( $n, d, \{v\}$ )  
... using class field theory **polsubcyclofast**( $n, d$ )  
roots of unity in  $nf$  **nfrootsof1**( $nf$ )  
roots of  $g$  belonging to  $nf$  **nfroots**( $nf, g$ )  
factor  $g$  in  $nf$  **nfactor**( $nf, g$ )

### Linear and algebraic relations

poly of degree  $\leq k$  with root  $x \in \mathbf{C}$  or  $\mathbf{Q}_p$  **algdep**( $x, k$ )  
alg. dep. with pol. coeffs for series  $s$  **seralgdep**( $s, x, y$ )  
diff. dep. with pol. coeffs for series  $s$  **serdiffdep**( $s, x, y$ )  
small linear rel. on coords of vector  $x$  **lindep**( $x$ )

## Basic Number Field Arithmetic (nf)

Number field elements are **t\_INT**, **t\_FRAC**, **t\_POL**, **t\_POLMOD**, or **t\_COL**  
(on integral basis  $nf.zk$ ).

### Basic operations

$x + y$  **nfeltadd**( $nf, x, y$ )  
 $x \times y$  **nfeltmul**( $nf, x, y$ )  
 $x^n$ ,  $n \in \mathbf{Z}$  **nfeltpow**( $nf, x, n$ )  
 $x/y$  **nfeltdiv**( $nf, x, y$ )  
 $q = x \setminus y := \text{round}(x/y)$  **nfeltdiveuc**( $nf, x, y$ )  
 $r = x \% y := x - (x \setminus y)y$  **nfeltmod**( $nf, x, y$ )  
...  $[q, r]$  as above **nfeltdivrem**( $nf, x, y$ )  
reduce  $x$  modulo ideal  $A$  **nfeltreduce**( $nf, x, A$ )  
absolute trace  $\text{Tr}_{K/\mathbf{Q}}(x)$  **nfelttrace**( $nf, x$ )  
absolute norm  $N_{K/\mathbf{Q}}(x)$  **nfeltnorm**( $nf, x$ )

is  $x$  a square? **nfeltissquare**( $nf, x, \{\&y\}$ )  
... an  $n$ -th power? **nfeltispower**( $nf, x, n, \{\&y\}$ )

**Multiplicative structure of  $K^*$ ;  $K^*/(K^*)^n$**   
valuation  $v_{\mathfrak{p}}(x)$  **nfeltval**( $nf, x, \mathfrak{p}$ )  
... write  $x = \pi^{v_{\mathfrak{p}}(x)}y$  **nfeltval**( $nf, x, \mathfrak{p}, \&y$ )  
quadratic Hilbert symbol (at  $\mathfrak{p}$ ) **nfhilbert**( $nf, a, b, \{\mathfrak{p}\}$ )  
 $b$  such that  $xb^n = v$  is small **idealredmodpower**( $nf, x, n$ )

### Maximal order and discriminant

integral basis of field **Q**[ $x$ ]/( $f$ ) **nfbasis**( $f$ )  
field discriminant of **Q**[ $x$ ]/( $f$ ) **nfdisc**( $f$ )  
... and factorization **nfdiscfactors**( $f$ )  
express  $x$  on integer basis **nfalgtobasis**( $nf, x$ )  
express element  $x$  as a polmod **nfbasistoalg**( $nf, x$ )

### Hecke Grossencharacters

Let  $K$  be a number field and  $m$  a modulus. A **gchar** structure describes the group of Hecke Grossencharacters of  $K$  of modulus  $m$  and allows computations with these characters. A character  $\chi$  is described by its components modulo  $gc.cyc$ .

init **gchar** structure  $gc$  for modulus  $m$  **gcharinit**( $bnf, m, \{cm\}$ )

### gc members:

  underlying  $bnf$  **gc.bnf**  
  modulus **gc.mod**  
  elementary divisors (including 0s) **gc.cyc**  
recompute  $gc$  using current precision **gcharnewprec**( $gc$ )  
evaluate Hecke character  $chi$  at ideal  $id$  **gchareval**( $gc, chi, id$ )  
exponent column of  $id$  in  $\mathbf{R}^n$  **gcharideallog**( $gc, id$ )  
log representation of ideal  $id$  **gcharlog**( $gc, id$ )  
... of character  $\chi$  **gcharduallog**( $gc, chi$ )  
exponent vector of  $\chi$  in  $\mathbf{R}^n$  **gcharparameters**( $gc, chi$ )  
conductor of  $\chi$  **gcharconductor**( $gc, chi$ )  
L-function of  $\chi$  **lfuncreate**( $[gc, chi]$ )  
local component  $\chi_v$  of  $\chi$  **gcharlocal**( $gc, chi, v$ )  
 $\chi$  s.t.  $\chi_v \approx Lchiv[i]$  for  $v = Lv[i]$  **gcharidentify**( $gc, Lv, Lchiv$ )  
basis of group of algebraic characters **gcharalgebraic**( $gc$ )  
is  $\chi$  algebraic? **gcharisalgebraic**( $gc, chi$ )

### Dedekind Zeta Function $\zeta_K$ , Hecke $L$ series

$R = [c, w, h]$  in initialization means we restrict  $s \in \mathbf{C}$  to domain  $|\Re(s) - c| < w$ ,  $|\Im(s)| < h$ ;  $R = [w, h]$  encodes  $[1/2, w, h]$  and  $[h]$  encodes  $R = [1/2, 0, h]$  (critical line up to height  $h$ ).

$\zeta_K$  as Dirichlet series,  $N(I) \leq b$  **dirzetak**( $nf, b$ )  
init  $\zeta_K^{(k)}(s)$  for  $k \leq n$  **L = lfunitinit**( $bnf, R, \{n = 0\}$ )  
compute  $\zeta_K(s)$  ( $n$ -th derivative) **lfun**( $L, s, \{n = 0\}$ )  
compute  $\Lambda_K(s)$  ( $n$ -th derivative) **lfunlambda**( $L, s, \{n = 0\}$ )

init  $L_K^{(k)}(s, \chi)$  for  $k \leq n$  **L = lfunitinit**( $[bnr, chi], R, \{n = 0\}$ )  
compute  $L_K(s, \chi)$  ( $n$ -th derivative) **lfun**( $L, s, \{n\}$ )  
Artin root number of  $K$  **bnrrootnumber**( $bnr, chi, \{flag\}$ )  
 $L(1, \chi)$ , for all  $\chi$  trivial on  $H$  **bnrL1**( $bnr, \{H\}, \{flag\}$ )

## Class Groups & Units (bnf, bnr)

Class field theory data  $a_1, \{a_2\}$  is usually  $bnr$  (ray class field),  $bnr, H$  (congruence subgroup) or  $bnr, \chi$  (character on **bnr.clgp**). Any of these define a unique abelian extension of  $K$ .  
units /  $S$ -units **bnfunits**( $bnf, \{S\}$ )  
remove GRH assumption from  $bnf$  **bnfcertify**( $bnf$ )

expo. of ideal $x$ on class gp	<code>bnfisprincipal(bnf, x, {flag})</code>
... on ray class gp	<code>bnrisprincipal(bnr, x, {flag})</code>
expo. of $x$ on fund. units	<code>bnfisunit(bnf, x)</code>
... on $S$ -units, $U$ is <code>bnfunits(bnf, S)</code>	<code>bnfisunit(bnfs, x, U)</code>
signs of real embeddings of $bnf$ .fu	<code>bnfsignunit(bnf)</code>
narrow class group	<code>bnfnarrow(bnf)</code>

### Class Field Theory

ray class number for modulus $m$	<code>bnrclassno(bnf, m)</code>
discriminant of class field	<code>bnrdisc(a1, {a2})</code>
ray class numbers, $l$ list of moduli	<code>bnrclassnolist(bnf, l)</code>
discriminants of class fields	<code>bnrdisclist(bnf, l, {arch}, {flag})</code>
decode output from <code>bnrdisclist</code>	<code>bnfdecodemodule(nf, fa)</code>
is modulus the conductor?	<code>bnrisconductor(a1, {a2})</code>
is class field $(bnr, H)$ Galois over $K^G$	<code>bnrisgalois(bnr, G, H)</code>
action of automorphism on <code>bnr.gen</code>	<code>bnrgaloismatrix(bnr, aut)</code>
apply <code>bnrgaloismatrix M</code> to $H$	<code>bnrgaloisapply(bnr, M, H)</code>
characters on <code>bnr.clgp</code> s.t. $\chi(g_i) = e(v_i)$	<code>bnrchar(bnr, g, {v})</code>
conductor of character $\chi$	<code>bnrconductor(bnr, chi)</code>
conductor of extension	<code>bnrconductor(a1, {a2}, {flag})</code>
conductor of extension $K[Y]/(g)$	<code>rnfconductor(bnf, g)</code>
canonical projection $\text{Cl}_F \rightarrow \text{Cl}_f$ , $f \mid F$	<code>bnrmap</code>
Artin group of extension $K[Y]/(g)$	<code>rnfnormgroup(bnr, g)</code>
subgroups of $bnr$ , index $\leq b$	<code>subgrouplist(bnr, b, {flag})</code>
compositum as <code>[bnr, H]</code>	<code>bnrcompositum([bnr1, H1], [bnr2, H2])</code>
class field defined by $H \subset \text{Cl}_f$	<code>bnrclassfield(bnr, H)</code>
... low level equivalent, prime degree	<code>rnfkummer(bnr, H)</code>
same, using Stark units (real field)	<code>bnrstark(bnr, sub, {flag})</code>
is $a$ an $n$ -th power in $K_v$ ?	<code>nfislocalpower(nf, v, a, n)</code>
cyclic $L/K$ satisf. local conditions	<code>nfgrunwaldwang(nf, P, D, pl)</code>

### Cyclotomic and Abelian fields theory

An Abelian field  $F$  given by a subgroup  $H \subset (Z/fZ)^*$  is described by an argument  $F$ , e.g.  $f$  (for  $H = 1$ , i.e.  $Q(\zeta_f)$ ) or  $[G, H]$ , where  $G$  is `idealstar(f, 1)`, or a minimal polynomial.

minus class number $h^-(F)$	<code>subcyclohminus(F)</code>
... $p$ -part	<code>subcyclohminus(F, p)</code>
minus part of Iwasawa polynomials	<code>subcycloiwasawa(F, p)</code>
$p$ -Sylog of $\text{Cl}(F)$	<code>subcyclopclgp(F, p)</code>

### Logarithmic class group

logarithmic $\ell$ -class group	<code>bnflog(bnf, l)</code>
$[\tilde{e}(F_v/Q_p), \tilde{f}(F_v/Q_p)]$	<code>bnflogef(bnf, pr)</code>
$\exp \deg_F(A)$	<code>bnflogdegree(bnf, A, l)</code>
is $\ell$ -extension $L/K$ locally cyclotomic	<code>rnfislocalcyclo(rnf)</code>

**Ideals:** elements, primes, or matrix of generators in HNF

is $id$ an ideal in $nf$ ?	<code>nfisideal(nf, id)</code>
is $x$ principal in $bnf$ ?	<code>bnfisprincipal(bnf, x)</code>
give $[a, b]$ , s.t. $a\mathbf{Z}_K + b\mathbf{Z}_K = x$	<code>idealtwoelt(nf, x, {a})</code>
put ideal $a$ ( $a\mathbf{Z}_K + b\mathbf{Z}_K$ ) in HNF form	<code>idealhnf(nf, a, {b})</code>
norm of ideal $x$	<code>idealnrm(nf, x)</code>
minimum of ideal $x$ (direction $v$ )	<code>idealmin(nf, x, v)</code>
LLL-reduce the ideal $x$ (direction $v$ )	<code>idealred(nf, x, {v})</code>

### Ideal Operations

add ideals $x$ and $y$	<code>idealadd(nf, x, y)</code>
multiply ideals $x$ and $y$	<code>idealmul(nf, x, y, {flag})</code>
intersection of ideal $x$ with $Q$	<code>idealdown(nf, x)</code>
intersection of ideals $x$ and $y$	<code>idealintersect(nf, x, y, {flag})</code>
$n$ -th power of ideal $x$	<code>idealpow(nf, x, n, {flag})</code>
inverse of ideal $x$	<code>idealinv(nf, x)</code>
divide ideal $x$ by $y$	<code>idealdiv(nf, x, y, {flag})</code>

# Algebraic Number Theory

(PARI-GP version 2.15.3)

Find $(a, b) \in x \times y$ , $a + b = 1$	<code>idealaddtoone(nf, x, {y})</code>
coprime integral $A, B$ such that $x = A/B$	<code>idealnumden(nf, x)</code>

### Primes and Multiplicative Structure

check whether $x$ is a maximal ideal	<code>idealismaximal(nf, x)</code>
factor ideal $x$ in $\mathbf{Z}_K$	<code>idealfactor(nf, x)</code>
expand ideal factorization in $K$	<code>idealfactorback(nf, f, {e})</code>
is ideal $A$ an $n$ -th power ?	<code>idealispower(nf, A, n)</code>
expand elt factorization in $K$	<code>nffactorback(nf, f, {e})</code>
decomposition of prime $p$ in $\mathbf{Z}_K$	<code>idealprimedec(nf, p)</code>
valuation of $x$ at prime ideal $pr$	<code>idealval(nf, x, pr)</code>
weak approximation theorem in $nf$	<code>idealchinese(nf, x, y)</code>
$a \in K$ , s.t. $v_p(a) = v_p(x)$ if $v_p(x) \neq 0$	<code>idealappr(nf, x)</code>
$a \in K$ such that $(a \cdot x, y) = 1$	<code>idealcoprime(nf, x, y)</code>
give $bid$ = structure of $(\mathbf{Z}_K/id)^*$	<code>idealstar(nf, id, {flag})</code>
structure of $(1 + \mathfrak{p})/(1 + \mathfrak{p}^k)$	<code>idealprincipalunits(nf, pr, k)</code>
discrete log of $x$ in $(\mathbf{Z}_K/bid)^*$	<code>ideallog(nf, x, bid)</code>
idealstar of all ideals of norm $\leq b$	<code>ideallist(nf, b, {flag})</code>
add Archimedean places	<code>ideallistarch(nf, b, {ar}, {flag})</code>
init <code>modpr</code> structure	<code>nfmodprinit(nf, pr, {v})</code>
project $t$ to $\mathbf{Z}_K/pr$	<code>nfmodpr(nf, t, modpr)</code>
lift from $\mathbf{Z}_K/pr$	<code>nfmodprlift(nf, t, modpr)</code>

### Galois theory over Q

conjugates of a root $\theta$ of $nf$	<code>nfgaloisconj(nf, {flag})</code>
apply Galois automorphism $s$ to $x$	<code>nfgaloisapply(nf, s, x)</code>
Galois group of field $\mathbf{Q}[x]/(f)$	<code>polgalois(f)</code>
resolvent field of $\mathbf{Q}[x]/(f)$	<code>nfresolvent(f)</code>
initializes a Galois group structure $G$	<code>galoisinit(pol, {den})</code>
... for the splitting field of $pol$	<code>galoisplittinginit(pol, {d})</code>
character table of $G$	<code>galoischartable(G)</code>
conjugacy classes of $G$	<code>galoisconjclasses(G)</code>
$\det(1 - \rho(g)T)$ , $\chi$ character of $\rho$	<code>galoischarpoly(G, x, {o})</code>
$\det(\rho(g))$ , $\chi$ character of $\rho$	<code>galoischarpoly(G, x, {o})</code>
action of $p$ in <code>nfgaloisconj</code> form	<code>galoispermtpol(G, {p})</code>
identify as abstract group	<code>galoisidentify(G)</code>
export a group for GAP/MAGMA	<code>galoisexport(G, {flag})</code>
subgroups of the Galois group $G$	<code>galoissubgroups(G)</code>
is subgroup $H$ normal?	<code>galoisisnormal(G, H)</code>
subfields from subgroups	<code>galoissubfields(G, {flag}, {v})</code>
fixed field	<code>galoisfixedfield(G, perm, {flag}, {v})</code>
Frobenius at maximal ideal $P$	<code>idealfrobenius(nf, G, P)</code>
ramification groups at $P$	<code>idealramgroups(nf, G, P)</code>
is $G$ abelian?	<code>galoisisabelian(G, {flag})</code>
abelian number fields/ $\mathbf{Q}$	<code>galoissubcyclo(N, H, {flag}, {v})</code>

### The galpol package

query the package: polynomial	<code>galoisgetpol(a, b, {s})</code>
...: permutation group	<code>galoisgetgroup(a, b)</code>
...: group description	<code>galoisgetname(a, b)</code>

# Relative Number Fields (rnf)

Extension  $L/K$  is defined by  $T \in K[x]$ .

absolute equation of $L$	<code>rnfequation(nf, T, {flag})</code>
is $L/K$ abelian?	<code>rnfisabelian(nf, T)</code>
relative <code>nfalgtobasis</code>	<code>rnfalgtobasis(rnf, x)</code>
relative <code>nfbasistoalg</code>	<code>rnfbasistoalg(rnf, x)</code>
relative <code>idealhnf</code>	<code>rnfidealhnf(rnf, x)</code>
relative <code>idealmul</code>	<code>rnfidealmul(rnf, x, y)</code>
relative <code>idealtwoelt</code>	<code>rnfidealtwoelt(rnf, x)</code>

### Lifts and Push-downs

absolute $\rightarrow$ relative representation for $x$	<code>rnfeltabstorel(rnf, x)</code>
relative $\rightarrow$ absolute representation for $x$	<code>rnfeltreltoabs(rnf, x)</code>
lift $x$ to the relative field	<code>rnfeltup(rnf, x)</code>
push $x$ down to the base field	<code>rnfeltdown(rnf, x)</code>
idem for $x$ ideal: <code>(rnfideal)reltoabs, abstorel, up, down</code>	

### Norms and Trace

relative norm of element $x \in L$	<code>rnfeltnrm(rnf, x)</code>
relative trace of element $x \in L$	<code>rnfelttrace(rnf, x)</code>
absolute norm of ideal $x$	<code>rnfidealnrmabs(rnf, x)</code>
relative norm of ideal $x$	<code>rnfidealnrmrel(rnf, x)</code>
solutions of $N_{K/\mathbf{Q}}(y) = x \in \mathbf{Z}$	<code>bnfisintnrm(bnf, x)</code>
is $x \in \mathbf{Q}$ a norm from $K$ ?	<code>bnfisnorm(bnf, x, {flag})</code>
initialize $T$ for norm eq. solver	<code>rnfisnorminit(K, pol, {flag})</code>
is $a \in K$ a norm from $L$ ?	<code>rnfisnorm(T, a, {flag})</code>
initialize $t$ for Thue equation solver	<code>thueinit(f)</code>
solve Thue equation $f(x, y) = a$	<code>thue(t, a, {sol})</code>
characteristic poly. of $a \bmod T$	<code>rnfcharpoly(nf, T, a, {v})</code>

### Factorization

factor ideal $x$ in $L$	<code>rnfidealfactor(rnf, x)</code>
$[S, T]: T_{i,j} \mid S_i$ ; $S$ primes of $K$ above $p$	<code>rnfidealprimedec(rnf, p)</code>

### Maximal order $\mathbf{Z}_L$ as a $\mathbf{Z}_K$ -module

relative <code>polredbest</code>	<code>rnfpolredbest(nf, T)</code>
relative <code>polredabs</code>	<code>rnfpolredabs(nf, T)</code>
relative Dedekind criterion, prime $pr$	<code>rnfdedekind(nf, T, pr)</code>
discriminant of relative extension	<code>rnfdisc(nf, T)</code>
pseudo-basis of $\mathbf{Z}_L$	<code>rnfpseudobasis(nf, T)</code>

**General  $\mathbf{Z}_K$ -modules:**  $M = [\text{matrix, vec. of ideals}] \subset L$

relative HNF / SNF	<code>nfhnf(nf, M), nfsnf</code>
multiple of det $M$	<code>nfdetint(nf, M)</code>
HNF of $M$ where $d = nfdetint(M)$	<code>nfhnfmod(x, d)</code>
reduced basis for $M$	<code>rnfilllgram(nf, T, M)</code>
determinant of pseudo-matrix $M$	<code>rnfdet(nf, M)</code>
Steinitz class of $M$	<code>rnfstteinitz(nf, M)</code>
$\mathbf{Z}_K$ -basis of $M$ if $\mathbf{Z}_K$ -free, or 0	<code>rnfhnfbasis(bnf, M)</code>
$n$ -basis of $M$ , or $(n + 1)$ -generating set	<code>rnfbasis(bnf, M)</code>
is $M$ a free $\mathbf{Z}_K$ -module?	<code>rnfisfree(bnf, M)</code>

Associative Algebras

$A$  is a general associative algebra given by a multiplication table  $mt$  (over  $\mathbf{Q}$  or  $\mathbf{F}_p$ ); represented by  $al$  from `algtableinit`.  
create  $al$  from  $mt$  (over  $\mathbf{F}_p$ )                    `algtableinit(mt, {p = 0})`  
group algebra  $\mathbf{Q}[G]$  (or  $\mathbf{F}_p[G]$ )                    `alggroup(G, {p = 0})`  
center of group algebra                    `alggrouppcenter(G, {p = 0})`  
**Properties**  
is  $(mt, p)$  OK for `algtableinit`?                    `algisassociative(mt, {p = 0})`  
multiplication table  $mt$                     `algmultable(al)`  
dimension of  $A$  over prime subfield                    `algdim(al)`  
characteristic of  $A$                     `algchar(al)`  
is  $A$  commutative?                    `algiscommutative(al)`  
is  $A$  simple?                    `algissimple(al)`  
is  $A$  semi-simple?                    `algissemisimple(al)`  
center of  $A$                     `algcenter(al)`  
Jacobson radical of  $A$                     `algradical(al)`  
radical  $J$  and simple factors of  $A/J$                     `algsimpledec(al)`  
**Operations on algebras**  
create  $A/I$ ,  $I$  two-sided ideal                    `algquotient(al, I)`  
create  $A_1 \otimes A_2$                     `algtensor(al1, al2)`  
create subalgebra from basis  $B$                     `algsubalg(al, B)`  
quotients by ortho. central idempotents  $e$                     `algcentralproj(al, e)`  
isomorphic alg. with integral mult. table                    `algmakeintegral(mt)`  
prime subalgebra of semi-simple  $A$  over  $\mathbf{F}_p$                     `algprimesubalg(al)`  
find isomorphism  $A \cong M_d(\mathbf{F}_q)$                     `algsplit(al)`  
**Operations on lattices in algebras**  
lattice generated by cols. of  $M$                     `alglathnf(al, M)`  
... by the products  $xy$ ,  $x \in lat1$ ,  $y \in lat2$                     `alglatmul(al, lat1, lat2)`  
sum  $lat1 + lat2$  of the lattices                    `alglatadd(al, lat1, lat2)`  
intersection  $lat1 \cap lat2$                     `alglatinter(al, lat1, lat2)`  
test  $lat1 \subset lat2$                     `alglatsubset(al, lat1, lat2)`  
generalized index  $(lat2 : lat1)$                     `alglatindex(al, lat1, lat2)`  
 $\{x \in al \mid x \cdot lat1 \subset lat2\}$                     `alglatlefttransporter(al, lat1, lat2)`  
 $\{x \in al \mid lat1 \cdot x \subset lat2\}$                     `alglatrighttransporter(al, lat1, lat2)`  
test  $x \in lat$  (set  $c = \text{coord. of } x$ )                    `alglatcontains(al, lat, x, {\&c})`  
element of  $lat$  with coordinates  $c$                     `alglatelement(al, lat, c)`  
**Operations on elements**  
 $a + b$ ,  $a - b$ ,  $-a$                     `algadd(al, a, b)`, `algsub`, `algneg`  
 $a \times b$ ,  $a^2$                     `algmul(al, a, b)`, `algsqr`  
 $a^n$ ,  $a^{-1}$                     `algpow(al, a, n)`, `alginv`  
is  $x$  invertible ? (then set  $z = x^{-1}$ )                    `algisinv(al, x, {\&z})`  
find  $z$  such that  $x \times z = y$                     `algdivl(al, x, y)`  
find  $z$  such that  $z \times x = y$                     `algdivr(al, x, y)`  
does  $z$  s.t.  $x \times z = y$  exist? (set it)                    `algisdivl(al, x, y, {\&z})`  
matrix of  $v \mapsto x \cdot v$                     `algtomatrix(al, x)`  
absolute norm                    `algnorm(al, x)`  
absolute trace                    `algtrace(al, x)`  
absolute char. polynomial                    `algcharpoly(al, x)`  
given  $a \in A$  and polynomial  $T$ , return  $T(a)$                     `algpoleval(al, T, a)`  
random element in a box                    `algrandom(al, b)`

Central Simple Algebras

$A$  is a central simple algebra over a number field  $K$ ; represented by  $al$  from `algininit`;  $K$  is given by a  $nf$  structure.  
create CSA from data                    `algininit(B, C, {v}, {maxord = 1})`  
multiplication table over  $K$                      $B = K$ ,  $C = mt$   
cyclic algebra  $(L/K, \sigma, b)$                      $B = rnf$ ,  $C = [sigma, b]$   
quaternion algebra  $(a, b)_K$                      $B = K$ ,  $C = [a, b]$   
matrix algebra  $M_d(K)$                      $B = K$ ,  $C = d$   
local Hasse invariants over  $K$                      $B = K$ ,  $C = [d, [PR, HF], HI]$

Properties

type of  $al$  ( $mt$ , CSA)                    `algtype(al)`  
dimension of  $A$  over  $\mathbf{Q}$                     `algdim(al, 1)`  
dimension of  $al$  over its center  $K$                     `algdim(al)`  
degree of  $A$  ( $= \sqrt{\dim_K A}$ )                    `algdegree(al)`  
 $al$  a cyclic algebra  $(L/K, \sigma, b)$ ; return  $\sigma$                     `algaut(al)`  
...return  $b$                     `algb(al)`  
...return  $L/K$ , as an  $rnf$                     `algsplittingfield(al)`  
split  $A$  over an extension of  $K$                     `algsplittingdata(al)`  
splitting field of  $A$  as an  $rnf$  over center                    `algsplittingfield(al)`  
multiplication table over center                    `algrelmultable(al)`  
places of  $K$  at which  $A$  ramifies                    `algramifiedplaces(al)`  
Hasse invariants at finite places of  $K$                     `alghassef(al)`  
Hasse invariants at infinite places of  $K$                     `alghassei(al)`  
Hasse invariant at place  $v$                     `alghasse(al, v)`  
index of  $A$  over  $K$  (at place  $v$ )                    `algindex(al, {v})`  
is  $al$  a division algebra? (at place  $v$ )                    `algisdivision(al, {v})`  
is  $A$  ramified? (at place  $v$ )                    `algisramified(al, {v})`  
is  $A$  split? (at place  $v$ )                    `algisplit(al, {v})`

Operations on elements

reduced norm                    `algnorm(al, x)`  
reduced trace                    `algtrace(al, x)`  
reduced char. polynomial                    `algcharpoly(al, x)`  
express  $x$  on integral basis                    `algalgtobasis(al, x)`  
convert  $x$  to algebraic form                    `algbasistoalg(al, x)`  
map  $x \in A$  to  $M_d(L)$ ,  $L$  split. field                    `algtomatrix(al, x)`

Orders

**Z**-basis of order  $\mathcal{O}_0$                     `algbasis(al)`  
discriminant of order  $\mathcal{O}_0$                     `algdisc(al)`  
**Z**-basis of natural order in terms  $\mathcal{O}_0$ 's basis                    `alginvbasis(al)`